

FORMULAS FOR POSITIVE, NEGATIVE AND ZERO VALUES OF THE MÖBIUS FUNCTION

R. M. ABRAROV¹ AND S. M. ABRAROV²

ABSTRACT. We obtained the formulas for the quantities of positive, negative and zero values of the Möbius function for any real x in terms of the Möbius function values for square root of x – similar to the identities we found earlier for the Mertens function [1]. Using the Dirac delta function approach [3, 2] we propose the equations showing how the Mertens and related functions can generate the output values.

Keywords: Möbius function, Mertens function, squarefree numbers, distribution of primes, Riemann Hypothesis.

1. Distributions of the Möbius function values

The Mertens function $M(x) = \sum_{k=1}^x \mu_k$ can be expressed through identity [1]

$$(1) \quad M(x) = 2M(\sqrt{x}) - \sum_{i,j=1}^{\sqrt{x}} \mu_i \mu_j \left[\frac{x}{i \cdot j} \right],$$

where μ_k is the Möbius function. It is possible to find similar identities for distributions of positive, negative and zero values of the Möbius function.

Let us denote $N_+(x)$, $N_-(x)$, $N_0(x)$ as the quantity of natural numbers correspondingly with $\mu = +1$, $\mu = -1$ and $\mu = 0$ in the interval not exceeding $x \geq 0$. For the Mertens function we have $M(x) = N_+(x) - N_-(x)$. We will also apply a notation $Q(x)$ [4, 6], commonly used for the quantity of squarefree numbers $N_+(x) + N_-(x)$.

There are obvious relations between quantities

$$(2) \quad \begin{aligned} [x] &= N_+(x) + N_-(x) + N_0(x), \\ 2N_+(x) &= [x] - N_0(x) + M(x), \\ 2N_-(x) &= [x] - N_0(x) - M(x). \end{aligned}$$

The frequencies of appearing numbers with $\mu = +1$, $\mu = -1$ and $\mu = 0$ are $\frac{N_+(x)}{[x]}$, $\frac{N_-(x)}{[x]}$ and $\frac{N_0(x)}{[x]}$, respectively. It should be noted

Date: May 4, 2009.

that the frequencies have some properties, similar to properties of the probabilities such as

$$(3) \quad \frac{N_+(x)}{[x]} + \frac{N_-(x)}{[x]} + \frac{N_0(x)}{[x]} = 1.$$

However we are considering here the deterministic process. Therefore these frequencies must not be confused with probabilities.

Let us find identities for $N_+(x)$, $N_-(x)$, $N_0(x)$ through known distribution of the Möbius function for $\sqrt{x}$, similar to (1). It is straightforward to get expression for $N_0(x)$ by using the *inclusion-exclusion principle* [7, 5]. In this case we count the quantity of numbers divisible by 2^2 plus numbers divisible by 3^2 , except those divisible by 2^2 and 3^2 , and so on for squares of all prime numbers:

$$(4) \quad \begin{aligned} N_0(x) = & \left[\frac{x}{2^2} \right] + \left[\frac{x}{3^2} \right] - \left[\frac{x}{2^2 \cdot 3^2} \right] \\ & + \left[\frac{x}{5^2} \right] - \left[\frac{x}{2^2 \cdot 5^2} \right] - \left[\frac{x}{3^2 \cdot 5^2} \right] + \left[\frac{x}{2^2 \cdot 3^2 \cdot 5^2} \right] + \dots \end{aligned}$$

If we rearrange the terms in order of the appearance of nonzero elements, we will get

$$(5) \quad N_0(x) = - \sum_{i=2}^{\sqrt{x}} \mu_i \left[\frac{x}{i^2} \right].$$

Hence from (5), (1) and (2) immediately follow

$$(6) \quad \begin{aligned} Q(x) &= \sum_{i=1}^{\sqrt{x}} \mu_i \left[\frac{x}{i^2} \right], \\ N_+(x) &= M(\sqrt{x}) - \frac{1}{2} \sum_{i,j=1}^{\sqrt{x}} \mu_i \mu_j \left[\frac{x}{i \cdot j} \right] + \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \mu_i \left[\frac{x}{i^2} \right], \\ N_-(x) &= -M(\sqrt{x}) + \frac{1}{2} \sum_{i,j=1}^{\sqrt{x}} \mu_i \mu_j \left[\frac{x}{i \cdot j} \right] + \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \mu_i \left[\frac{x}{i^2} \right]. \end{aligned}$$

2. Dirac and Kronecker delta function transformation

Formula (1) involves the floor function and explicit dependence on the Mertens function in the right hand side. In order to avoid them, we can apply an alternative for the floor function - the Dirac delta function δ^D [3] (see also our [2]) or the discrete version of the Dirac delta function $\delta^K(x) \equiv \delta_{[x][x]}$ - the Kronecker delta with floor and

ceiling functions of x as arguments (which means that the value of $\delta^K(x)$ is 1 only for integer x and 0 otherwise)

$$(7) \quad \left[\frac{x}{i \cdot j} \right] = \int_{i \cdot j - \varepsilon}^x \delta^D \left(\frac{y}{i \cdot j} - \left[\frac{y}{i \cdot j} \right] \right) dy$$

$$= \int_{1-\varepsilon}^x \delta^D \left(\frac{y}{i \cdot j} - \left[\frac{y}{i \cdot j} \right] \right) dy, \quad \text{here and further } x \geq 1,$$

$$(8) \quad \left[\frac{x}{i \cdot j} \right] = \sum_{k=i \cdot j}^x \delta^K \left(\frac{k}{i \cdot j} \right) = \sum_{k=1}^x \delta^K \left(\frac{k}{i \cdot j} \right).$$

Now substituting (7) and (8) into (1), we derive

$$(9) \quad M(x) = 2 - \int_{1^2 - \varepsilon}^{x \geq 1^2} \mu_1 \left(\mu_1 \delta^D \left(\frac{y}{1 \cdot 1} - \left[\frac{y}{1 \cdot 1} \right] \right) \right) dy$$

$$- \int_{2^2 - \varepsilon}^{x \geq 2^2} \mu_2 \left(2\mu_1 \delta^D \left(\frac{y}{2 \cdot 1} - \left[\frac{y}{2 \cdot 1} \right] \right) \right.$$

$$+ \left. \mu_2 \delta^D \left(\frac{y}{2 \cdot 2} - \left[\frac{y}{2 \cdot 2} \right] \right) \right) dy$$

$$- \int_{3^2 - \varepsilon}^{x \geq 3^2} \mu_3 \left(2\mu_1 \delta^D \left(\frac{y}{3 \cdot 1} - \left[\frac{y}{3 \cdot 1} \right] \right) \right.$$

$$+ \left. 2\mu_2 \delta^D \left(\frac{y}{3 \cdot 2} - \left[\frac{y}{3 \cdot 2} \right] \right) + \mu_3 \delta^D \left(\frac{y}{3 \cdot 3} - \left[\frac{y}{3 \cdot 3} \right] \right) \right) dy$$

$$- \int_{5^2 - \varepsilon}^{x \geq 5^2} \mu_5 \left(2\mu_1 \delta^D \left(\frac{y}{5 \cdot 1} - \left[\frac{y}{5 \cdot 1} \right] \right) \right.$$

$$+ \left. 2\mu_2 \delta^D \left(\frac{y}{5 \cdot 2} - \left[\frac{y}{5 \cdot 2} \right] \right) + 2\mu_3 \delta^D \left(\frac{y}{5 \cdot 3} - \left[\frac{y}{5 \cdot 3} \right] \right) \right.$$

$$+ \left. \mu_5 \delta^D \left(\frac{y}{5 \cdot 5} - \left[\frac{y}{5 \cdot 5} \right] \right) \right) dy - \dots$$

$$= 2 - \sum_{i=1}^{\sqrt{x}} \int_{i^2 - \varepsilon}^x \mu_i \left(-\mu_i \delta^D \left(\frac{y}{i \cdot i} - \left[\frac{y}{i \cdot i} \right] \right) \right.$$

$$+ \left. \sum_{j=1}^i 2\mu_j \delta^D \left(\frac{y}{i \cdot j} - \left[\frac{y}{i \cdot j} \right] \right) \right) dy,$$

$$\begin{aligned}
(10) \quad M(x) &= 2 - \sum_{k=1}^x \sum_{i,j=1}^{\sqrt{k}} \mu_i \mu_j \delta^K \left(\frac{k}{i \cdot j} \right) \\
&= 2 - \sum_{i=1}^{\sqrt{x}} \sum_{k=i^2}^x \mu_i \left(-\mu_i \delta^K \left(\frac{k}{i \cdot i} \right) + \sum_{j=1}^i 2\mu_j \delta^K \left(\frac{k}{i \cdot j} \right) \right).
\end{aligned}$$

From (10) we have

$$(11) \quad \mu_x = - \sum_{i,j=1}^{\sqrt{x}} \mu_i \mu_j \delta^K \left(\frac{x}{i \cdot j} \right) \quad x \geq 2.$$

By following the same procedure as it was in the previous section, we obtain

$$\begin{aligned}
(12) \quad N_0(x) &= - \sum_{i=2}^{\sqrt{x}} \mu_i \int_{i^2-\varepsilon}^x \delta^D \left(\frac{y}{i \cdot i} - \left[\frac{y}{i \cdot i} \right] \right) dy \\
&= - \sum_{i=2}^{\sqrt{x}} \sum_{k=i^2}^x \mu_i \delta^K \left(\frac{k}{i \cdot i} \right),
\end{aligned}$$

$$(13) \quad Q(x) = \sum_{i=1}^{\sqrt{x}} \mu_i \int_{i^2-\varepsilon}^x \delta^D \left(\frac{y}{i \cdot i} - \left[\frac{y}{i \cdot i} \right] \right) dy = \sum_{i=1}^{\sqrt{x}} \sum_{k=i^2}^x \mu_i \delta^K \left(\frac{k}{i \cdot i} \right),$$

$$\begin{aligned}
(14) \quad N_+(x) &= 1 - \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \int_{i^2-\varepsilon}^x \mu_i \left(-\mu_i \delta^D \left(\frac{y}{i \cdot i} - \left[\frac{y}{i \cdot i} \right] \right) \right. \\
&\quad \left. + \sum_{j=1}^i 2\mu_j \delta^D \left(\frac{y}{i \cdot j} - \left[\frac{y}{i \cdot j} \right] \right) \right) dy \\
&\quad + \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \mu_i \int_{i^2-\varepsilon}^x \delta^D \left(\frac{y}{i \cdot i} - \left[\frac{y}{i \cdot i} \right] \right) dy \\
&= 1 - \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \sum_{k=i^2}^x \mu_i \left(-\mu_i \delta^K \left(\frac{k}{i \cdot i} \right) \right. \\
&\quad \left. + \sum_{j=1}^i 2\mu_j \delta^K \left(\frac{k}{i \cdot j} \right) \right) + \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \sum_{k=i^2}^x \mu_i \delta^K \left(\frac{k}{i \cdot i} \right),
\end{aligned}$$

$$\begin{aligned}
 (15) \quad N_-(x) &= -1 + \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \int_{i^2-\varepsilon}^x \mu_i \left(-\mu_i \delta^D \left(\frac{y}{i \cdot i} - \left[\frac{y}{i \cdot i} \right] \right) \right. \\
 &\quad \left. + \sum_{j=1}^i 2\mu_j \delta^D \left(\frac{y}{i \cdot j} - \left[\frac{y}{i \cdot j} \right] \right) \right) dy \\
 &\quad + \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \mu_i \int_{i^2-\varepsilon}^x \delta^D \left(\frac{y}{i \cdot i} - \left[\frac{y}{i \cdot i} \right] \right) dy \\
 &= -1 + \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \sum_{k=i^2}^x \mu_i \left(-\mu_i \delta^K \left(\frac{k}{i \cdot i} \right) \right. \\
 &\quad \left. + \sum_{j=1}^i 2\mu_j \delta^K \left(\frac{k}{i \cdot j} \right) \right) + \frac{1}{2} \sum_{i=1}^{\sqrt{x}} \sum_{k=i^2}^x \mu_i \delta^K \left(\frac{k}{i \cdot i} \right).
 \end{aligned}$$

These identities are showing a clear mechanism how functions above generate the output values and can be helpful in understanding of their distributions and other properties.

REFERENCES

- [1] R. M. Abrarov and S. M. Abrarov, *Riemann Hypothesis may be proved by induction*, http://arxiv.org/PS_cache/arxiv/pdf/0802/0802.1764v1.pdf
- [2] S. M. Abrarov and R. M. Abrarov (submitted to journal), *Identities for number series and their reciprocals: Dirac delta function approach*, <http://arxiv.org/ftp/arxiv/papers/0704/0704.1936.pdf>
- [3] P. Antosik, J. Mikusinski, R. Sikorski, *Theory of Distributions. The Sequential Approach*, Elsevier Scientific Publishing Company, 1973.
- [4] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 5th ed., Oxford University Press, Oxford, 1979.
- [5] Havil, J. *Gamma: Exploring Euler's Constant*. Princeton, NJ: Princeton University Press, p. 66, 2003.
- [6] Weisstein, Eric W. "Squarefree." From MathWorld - A Wolfram Resource. <http://mathworld.wolfram.com/Squarefree.html>
- [7] Weisstein, Eric W. "Inclusion-Exclusion Principle." From MathWorld - A Wolfram Resource. <http://mathworld.wolfram.com/Inclusion-ExclusionPrinciple.html>

E-mail address: ¹rabrarov@gmail.com

E-mail address: ²absanj@gmail.com